

p -elementary subgroups of the Cremona group

Arnaud BEAUVILLE

Introduction

Let k be an algebraically closed field. The *Cremona group* Cr_k is the group of birational transformations of $\mathbb{P}_k^2$, or equivalently the group of k -automorphisms of the field $k(x, y)$. There is an extensive classical literature about this group, in particular about its finite subgroups – see the introduction of [dF] for a list of references.

The classification of conjugacy classes of elements of prime order p in Cr_k has been given a modern treatment in [B-B] for $p = 2$ and in [dF] for $p \geq 3$ (see also [B-B]). In this note we go one step further and classify p -elementary subgroups – that is, subgroups isomorphic to $(\mathbb{Z}/p)^r$ for p prime. We will mostly describe such a subgroup as a group G of automorphisms of a rational surface S : we identify G to a subgroup of Cr_k by choosing a birational map $\varphi : S \dashrightarrow \mathbb{P}^2$. Then the conjugacy class of G in Cr_k depends only on the data (G, S) .

Theorem. – *Let G be a subgroup of Cr_k of the form $(\mathbb{Z}/p)^r$ with p prime $\neq \mathrm{char}(k)$. Then:*

a) *Assume $p \geq 5$. Then $r \leq 2$, and if $r = 2$ G is conjugate to the p -torsion subgroup of the diagonal torus¹ of $\mathrm{PGL}_3(k) = \mathrm{Aut}(\mathbb{P}^2)$.*

b) *Assume $p = 3$. Then $r \leq 3$, and if $r = 3$ G is conjugate to the 3-torsion subgroup of the diagonal torus of $\mathrm{PGL}_4(k)$, acting on the Fermat cubic surface $X_0^3 + \dots + X_3^3 = 0$.*

c) *Assume $p = 2$. Then $r \leq 4$, and if equality holds G is conjugate to one of the following subgroups:*

c₁) *the subgroup of Cr_k spanned by the involutions:*

$$(x, y) \mapsto (-x, y) \quad , \quad (x, y) \mapsto \left(\frac{1}{x}, y\right) \quad , \quad (x, y) \mapsto (x, -y) \quad , \quad (x, y) \mapsto \left(x, \frac{P(x)}{y}\right)$$

with $P(x) = \prod_{a \in I} (x^2 + x^{-2} - a)$ for some finite subset I of $k \setminus \{2, -2\}$.

c₂) *the 2-torsion subgroup of the diagonal torus of $\mathrm{PGL}_5(k)$, acting on the quartic del Pezzo surface in $\mathbb{P}^4$ with equations $\sum_{i=0}^4 X_i^2 = \sum_{i=0}^4 \lambda_i X_i^2 = 0$ for some distinct elements $\lambda_0, \dots, \lambda_4$ of k .*

¹ By the *diagonal torus* of $\mathrm{PGL}_r(k)$ we mean the subgroup of projective transformations $(X_0, \dots, X_r) \mapsto (t_0 X_0, \dots, t_r X_r)$ for $t_0, \dots, t_r$ in k^* .

Note that the question does not make sense when $p = \text{char}(k)$, already in dimension 1: the group $\text{PGL}_2(k)$ contains a subgroup isomorphic to k , hence infinite-dimensional over $\mathbb{Z}/p$.

In the next section we discuss some motivation for this question; then we reduce the problem through standard techniques to the study of p -elementary subgroups $G \subset \text{Aut}(S)$, where S is either a del Pezzo surface or carries a $\mathbb{P}^1$ -fibration preserved by G . We will study the latter case in § 2 and the former in § 3. Finally in § 4 we discuss the classification of the conjugacy classes of subgroups isomorphic to $(\mathbb{Z}/2)^4$ (the only case where the conjugacy class is not unique).

As I. Dolgachev pointed out to us, the result (over $\mathbb{C}$) could be deduced from the list of the finite subgroups of the Cremona group established by Kantor [K], and completed by Wiman [W]. However, the results of § 2 and § 4 would still be needed to decide whether certain subgroups are conjugate or not. Most of the results of § 3 are contained in those of Kantor and Wiman, but they are so much simpler in our specific situation that we have preferred to give an independent proof.

I am indebted to J.-P. Serre for asking me the question considered here, and for useful discussions, in particular about (1.1) and (1.2) below. I thank the referee for pointing out a gap in the first version of this note.

1. Comments, and beginning of the proof

(1.1) Though very large, the Cremona group behaves in some respect like a semi-simple group of rank 2: every maximal torus has dimension 2, and is conjugate to the diagonal torus T of $\text{PGL}_3(k)$ [D1]. In this set-up the analogue of the Weyl group is the whole automorphisms group $\text{GL}_2(\mathbb{Z})$ of T ([D1], cor. 5 p. 522).

Now let H be a semi-simple group over k , and p a prime number ≥ 7 which does not divide the order of $\pi_1(H)$. Then every maximal p -elementary subgroup $G \subset H$ is the p -torsion subgroup of a maximal torus of H ; moreover this torus is unique, in fact it is the centralizer of G in H [Bo].

(1.2) Our theorem shows that the first part of this statement also holds for the Cremona group (with $p \geq 5$). However, *the maximal torus containing G is not unique*. Indeed, let T be the diagonal torus of $\text{PGL}_3(k)$, and G its p -torsion subgroup. The centralizer of G in Cr_k contains the transformations $\sigma_f : (x, y) \mapsto (x, yf(x^p))$ for $f \in k(t)^*$. If f is not a monomial σ_f does not normalize T , and G is also contained in $\sigma_f T \sigma_f^{-1}$.

(1.3) Now let us begin the proof of the Theorem. Let G be a finite subgroup of Cr_k . Then G can be realized as a group of automorphisms of a rational surface S (see for instance [dF-E], Thm. 1.4). Moreover we can assume that (G, S) is minimal,

that is, every birational G -equivariant morphism of S onto a smooth surface with a G -action is an isomorphism. Then one of the following holds:

- G preserves a fibration $f : S \rightarrow \mathbb{P}^1$ with rational fibers;
- $\text{rk Pic}(S)^G = 1$.

This result goes back to Manin [M], at least in the case (of interest for us) when G is abelian. It is by now a direct consequence of Mori theory, see for instance [Z], lemma 4.3.

In the former case G embeds in the group of automorphisms of the generic fibre $\mathbb{P}_{k(t)}^1$ of f ; in the next section we are going to classify the p -elementary subgroups of $\text{Aut}(\mathbb{P}_{k(t)}^1)$. In the latter case S is a del Pezzo surface, and the group $\text{Aut}(S)$ is well-known; we will use this information in § 3 to classify the corresponding p -elementary subgroups. Putting these results together gives the Theorem.

2. Subgroups of $\text{Aut}(\mathbb{P}_K^1)$

Let K be an extension of k , and p a prime number $\neq \text{char}(k)$. Let us recall the classification of p -elementary subgroups of $\text{PGL}_2(K)$. Let $C_p \subset \text{PGL}_2(K)$ be the cyclic subgroup of homographies $z \mapsto \zeta z$, with $\zeta^p = 1$; for $\delta \in K^*$, let $V_\delta \subset \text{PGL}_2(K)$ be the subgroup of order 4 spanned by the homographies $z \mapsto -z$ and $z \mapsto \frac{\delta}{z}$.

Lemma 2.1.— *Let G be a subgroup of $\text{PGL}_2(K)$ of the form $(\mathbb{Z}/p)^r$, with p prime $\neq \text{char}(k)$. Then G is conjugate to a subgroup of C_p if p is odd, and to a subgroup of V_δ for some $\delta \in K^*$ if $p = 2$.*

Proof: Let σ be an element of G , represented by a matrix $A \in \text{GL}_2(K)$ which satisfies $A^p = \delta I$ for some scalar $\delta \in K^*$. Taking determinants give $(\det A)^p = \delta^2$, so that $\delta^2 \equiv 1 \pmod{K^{*p}}$. If p is odd, this implies $\delta \in K^{*p}$; thus A is diagonalizable, and σ is conjugate to an element of C_p . The centralizer of C_p in $\text{PGL}_2(K)$ is the group of homotheties $z \mapsto \lambda z$, hence G is contained in its p -torsion subgroup C_p .

We now consider the case $p = 2$. In a basis of K^2 of the form (v, Av) , we have $A = \begin{pmatrix} 0 & \delta \\ 1 & 0 \end{pmatrix}$, so that σ becomes the homography $z \mapsto \frac{\delta}{z}$. The centralizer of σ in $\text{PGL}_2(\overline{K})$ is isomorphic to $\mathbb{Z}/2 \times \overline{K}^*$, so its 2-torsion subgroup is isomorphic to $(\mathbb{Z}/2)^2$; since this subgroup contains V_δ , it is equal to V_δ , and therefore $G \subset V_\delta$. ■

The conjugacy class of the subgroup V_δ depends only on the class of δ in K^*/K^{*2} . In particular, when $K = k$, all these subgroups are conjugate to V_1 .

(2.2) We are interested in the automorphism group $\text{Aut}(\mathbb{P}_K^1)$ of the k -scheme $\mathbb{P}_K^1$. Let Γ be the automorphism group of K over k . The action of $\text{Aut}(\mathbb{P}_K^1)$ on

the global functions of $\mathbb{P}_K^1$ gives rise to an exact sequence

$$1 \rightarrow \mathrm{PGL}_2(K) \longrightarrow \mathrm{Aut}(\mathbb{P}_K^1) \xrightarrow{\pi} \Gamma \rightarrow 1 ;$$

the surjection π has a canonical section $s : \Gamma \rightarrow \mathrm{Aut}(\mathbb{P}_K^1)$ which maps $\sigma \in \Gamma$ to the automorphism $z \mapsto \sigma(z)$. We will use this section to identify Γ to a subgroup of $\mathrm{Aut}(\mathbb{P}_K^1)$, and view $\mathrm{Aut}(\mathbb{P}_K^1)$ as the semi-direct product $\mathrm{PGL}_2(K) \rtimes \Gamma$. In other words, an element of $\mathrm{Aut}(\mathbb{P}_K^1)$ is a transformation $z \mapsto \frac{a\sigma(z) + b}{c\sigma(z) + d}$ for some

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(K) \text{ and } \sigma \in \Gamma.$$

We are interested in the case $K = k(t)$, so that $\Gamma = \mathrm{PGL}_2(k)$.

Proposition 2.3.— *Let G be a subgroup of $\mathrm{Aut}(\mathbb{P}_{k(t)}^1)$ of the form $(\mathbb{Z}/p)^r$, with p prime $\neq \mathrm{char}(k)$.*

a) *When p is odd, we have $r \leq 2$; if $r = 2$, G is conjugate to the subgroup $C_p \times C_p$ of $\mathrm{PGL}_2(K) \rtimes \mathrm{PGL}_2(k)$.*

b) *When $p = 2$ we have $r \leq 4$; if equality holds, G is conjugate to the subgroup $V_\delta \times V_1$ of $\mathrm{PGL}_2(K) \rtimes \mathrm{PGL}_2(k)$ for some element δ of K^* invariant under V_1 .*

Proof: Let $G' = G \cap \mathrm{PGL}_2(K)$. The bound on r follows from the exact sequence

$$1 \rightarrow G' \longrightarrow G \xrightarrow{\pi} \pi(G) \rightarrow 1$$

and Lemma 2.1. Assume that p is odd and $r = 2$. Then G' and $\pi(G)$ are cyclic, and after conjugation we may assume $G' = C_p$ and $\pi(G) = C_p$. Let σ be a generator of C_p , and g an element of G with $\pi(g) = \sigma$. We have $g = h\sigma$ for some element h of $\mathrm{PGL}_2(K)$. Since the elements of G' commute with σ and $h\sigma$, they must commute with h ; thus h belongs to the centralizer of $G' = C_p$ in $\mathrm{PGL}_2(K)$, hence is a homothety $z \mapsto \lambda z$ for some $\lambda \in K^*$. The relation $(h\sigma)^p = 1$ gives $\lambda \cdot \sigma(\lambda) \cdot \dots \cdot \sigma^{p-1}(\lambda) = 1$; by Hilbert's theorem 90, λ is then equal to $\mu^{-1}\sigma(\mu)$ for some $\mu \in K^*$. The homothety $z \mapsto \mu z$ conjugates g to σ , and commutes with G' , hence conjugates G to $C_p \times C_p$.

Now we assume $p = 2$ and $r = 4$. After conjugation we may assume $\pi(G) = V_1$, and $G' = V_\zeta$ for some $\zeta \in K^*$ (Lemma 2.1). Let $\sigma \in V_1$. The elements of $\pi^{-1}(\sigma)$ are of the form $h\sigma$ for some h in $\mathrm{PGL}_2(K)$. Since h must commute with the homography $z \mapsto -z$, it is of the form $z \mapsto \lambda z$ or $z \mapsto \frac{\mu}{z}$. Imposing that $h\sigma$ commutes with the homography $z \mapsto \frac{\zeta}{z}$ gives $\pi^{-1}(\sigma) = G' \cdot g_\sigma$, where $g_\sigma(z) = \lambda_\sigma \sigma(z)$, and $\lambda_\sigma \in K^*$ is a square root of $\zeta \sigma(\zeta)^{-1}$. Note that λ_σ is well-defined up to sign.

We can choose each λ_σ so that $\sigma \mapsto g_\sigma$ is a section of π (fix λ_σ arbitrarily on a basis of V_1 over $\mathbb{Z}/2$, and extend linearly). Then $\sigma \mapsto \lambda_\sigma$ is a 1-cocycle of V_1

with values in K^* . As above this cocycle is a coboundary, hence there exists $\mu \in K^*$ such that $\lambda_\sigma = \mu^{-1}\sigma(\mu)$ for each $\sigma \in V_1$. The homothety $z \mapsto \mu z$ conjugates g_σ to σ and V_ζ to V_δ , with $\delta = \zeta\mu^2$. The commutativity of G imposes that δ is invariant under V_1 ; this implies in particular that V_1 acts trivially on V_δ , hence $G = V_\delta \times V_1 \subset \mathrm{PGL}_2(K) \rtimes \mathrm{PGL}_2(k)$. ■

(2.4) Thus for p odd and $r = 2$, G is conjugate to the group of diagonal transformations $(z, t) \mapsto (\alpha z, \beta t)$, with $\alpha, \beta \in \mu_p$. For $p = 2$ and $r = 4$, G is conjugate to the group $V_\delta \times V_1$ generated by

$$(z, t) \mapsto (z, -t) \quad , \quad (z, t) \mapsto (z, \frac{1}{t}) \quad , \quad (z, t) \mapsto (-z, t) \quad , \quad (z, t) \mapsto (\frac{\delta}{z}, t) .$$

Since the map $t \mapsto t^2 + t^{-2}$ identifies $\mathbb{P}^1/V_1$ with $\mathbb{P}^1$, the invariance of δ under V_1 means that δ can be written $Q(t^2 + t^{-2})$ for some $Q \in k(s)$. Replacing Q by QF^2 does not change the conjugacy class of $V_\delta \times V_1$, so we can assume that Q is a polynomial with simple roots. Moreover conjugation by the birational transformation $(z, t) \mapsto (z(t \pm t^{-1}), t)$ amounts to multiply Q by $(s \pm 2)$, so we can assume that Q does not vanish at ± 2 . This gives case c_1) of the Theorem.

3. Automorphisms of del Pezzo surfaces

We now consider the case where S is a del Pezzo surface and $G \cong (\mathbb{Z}/p)^r$ a subgroup of $\mathrm{Aut}(S)$ such that $\mathrm{rk} \mathrm{Pic}(S)^G = 1$. We first recall the following well-known fact, which is a particular case of the results mentioned in (1.1):

Lemma 3.1. — *Let G be a subgroup of $\mathrm{PGL}_n(k)$ of the form $(\mathbb{Z}/p)^r$, with p prime $\neq \mathrm{char}(k)$. Assume that p does not divide n . Then $r \leq n - 1$, and if equality holds G is conjugate to the p -torsion subgroup of the diagonal torus.*

Proof : Pulling back G to $\mathrm{SL}_n(k)$ gives a central extension of G by the group μ_n of n -th roots of unity in k . Such extensions are parametrized by the group $H^2(G, \mu_n)$ which is annihilated both by n and p . Thus our extension is trivial, and G lifts to a subgroup of $\mathrm{SL}_n(k)$, isomorphic to $(\mathbb{Z}/p)^r$. Such a subgroup is contained in a maximal torus of $\mathrm{SL}_n(k)$, hence our assertions. ■

(3.2) Let us start with the case $S = \mathbb{P}^2$. By the above lemma, if $p \neq 3$, we have $r \leq 2$, and any subgroup of $\mathrm{PGL}_3(k)$ isomorphic to $(\mathbb{Z}/p)^2$ is conjugate to the p -torsion subgroup of the diagonal torus.

The case $p = 3$ is classical (see e.g. [Bo], 6.4 for a more general statement): we have again $r \leq 2$, and a subgroup isomorphic to $(\mathbb{Z}/3)^2$ is conjugate either to the diagonal subgroup, or to the subgroup spanned by the automorphisms $(X_0, X_1, X_2) \mapsto (X_1, X_2, X_0)$ and $(X_0, X_1, X_2) \mapsto (X_0, \alpha X_1, \alpha^2 X_2)$ for $\alpha \in \mu_3$.

(3.3) If S is obtained from $\mathbb{P}^2$ by blowing up one or two points, the group $\text{Aut}(S)$ is a subgroup of $\text{PGL}_3(k)$, so (3.2) applies. Suppose that S is the blow up of $\mathbb{P}^2$ at three non-collinear points. Let $N \subset \text{PGL}_3(k)$ be the subgroup of automorphisms preserving this three-points subset. The group $\text{Aut}(S)$ is the semi-direct product of N and a subgroup of order 2. Let G be a subgroup of $\text{Aut}(S)$ isomorphic to $(\mathbb{Z}/p)^r$. If $p \neq 2$, G is contained in N , so that (3.2) applies. If $p = 2$, we have $r \leq 3$ again by (3.2).

We now consider the case $S = \mathbb{P}^1 \times \mathbb{P}^1$.

Lemma 3.4.— *Let G be a group of automorphisms of $\mathbb{P}^1 \times \mathbb{P}^1$, isomorphic to $(\mathbb{Z}/p)^r$, such that $\text{rk Pic}(\mathbb{P}^1 \times \mathbb{P}^1)^G = 1$. Then $p = 2$ and $r \leq 3$.*

Proof : The automorphism group of $\mathbb{P}^1 \times \mathbb{P}^1$ is the semi-direct product $(\text{PGL}_2(k) \times \text{PGL}_2(k)) \rtimes \mathbb{Z}/2$, where $\mathbb{Z}/2$ acts on $\text{PGL}_2(k) \times \text{PGL}_2(k)$ by exchanging the factors. If $p \neq 2$, our subgroup G is contained in $\text{PGL}_2(k) \times \text{PGL}_2(k)$, hence $\text{Pic}(\mathbb{P}^1 \times \mathbb{P}^1)^G = \text{Pic}(\mathbb{P}^1 \times \mathbb{P}^1)$ has rank 2.

Thus we have $p = 2$. The subgroup G' of G preserving the two $\mathbb{P}^1$ -fibrations is contained in $\text{PGL}_2(k) \times \text{PGL}_2(k)$, thus in $C_2 \times C_2$ up to conjugacy, and G is conjugate to a subgroup of the semi-direct product $(C_2 \times C_2) \rtimes \mathbb{Z}/2$. But the elements of order 2 in this group not contained in $C_2 \times C_2$ are contained in the (direct) product of $\mathbb{Z}/2$ by the diagonal subgroup $C_2 \subset C_2 \times C_2$. Therefore G must be contained in $C_2 \times \mathbb{Z}/2$, hence $r \leq 3$. ■

(3.5) It remains to consider the case when S is obtained from $\mathbb{P}^2$ by blowing up ℓ points in general position, with $4 \leq \ell \leq 8$. We start by recalling some classical facts about such surfaces, which can be found for instance in [D2]. The primitive cohomology $H^2(S, \mathbb{Z})_{\text{prim}}$ (the orthogonal of the canonical class in $H^2(S, \mathbb{Z})$) is the root lattice of a root system R . The group $\text{Aut}(S)$ acts faithfully on $H^2(S, \mathbb{Z})_{\text{prim}}$, hence can be identified with a subgroup of the automorphism group of the root system R ; it is actually contained in the Weyl group $W \subset \text{Aut}(R)$ [Do]. The root systems which appear are the following (see [B]):

- $\ell = 4$: $R = A_4$, $W = \mathfrak{S}_5$
- $\ell = 5$: $R = D_5$, $W = (\mathbb{Z}/2)^4 \rtimes \mathfrak{S}_5$
- $\ell = 6$: $R = E_6$, $|W| = 2^7 \cdot 3^4 \cdot 5$
- $\ell = 7$: $R = E_7$, $|W| = 2^{10} \cdot 3^4 \cdot 5 \cdot 7$
- $\ell = 8$: $R = E_8$, $|W| = 2^{14} \cdot 3^5 \cdot 5^2 \cdot 7$

(3.6) For $\ell = 7$ (resp. $\ell = 8$), the linear system $| -K_S |$ (resp. $| -2K_S |$) defines a degree 2 morphism onto $\mathbb{P}^2$ (resp. a quadric cone in $\mathbb{P}^3$), branched along a canonically embedded smooth curve C of genus 3 (resp. 4). This gives rise to an

exact sequence

$$1 \rightarrow \mathbb{Z}/2 \longrightarrow \text{Aut}(S) \longrightarrow \text{Aut}(C) \quad (3.6)$$

(the right hand side map is actually surjective, but we will not need this). To control subgroups of $\text{Aut}(C)$ the following lemma will be useful:

Lemma 3.7.— *Let p be a prime number, r an integer, and C a curve of genus g with a faithful action of the group $(\mathbb{Z}/p)^r$. Then p^{r-1} divides $2g - 2$.*

Proof: Put $G = (\mathbb{Z}/p)^r$, and consider the covering $\pi : C \rightarrow C/G$. Since the stabilizer of any point of C is cyclic, the fibre of π above a branch point consists of p^{r-1} points with ramification index p . Thus Hurwitz's formula gives the result. ■

Proposition 3.8.— *For p prime ≥ 5 , the group $(\mathbb{Z}/p)^2$ does not act faithfully on a del Pezzo surface.*

Proof: A glance at the list 3.5 shows that the only case where the order of the Weyl group is divisible by p^2 , with $p \geq 5$, is $\ell = 8$. In this case the exact sequence (3.6) shows that $(\mathbb{Z}/p)^2$ acts faithfully on a smooth curve C of genus 4, and this contradicts Lemma 3.7.

Proposition 3.9.— *Let S be a del Pezzo surface admitting a faithful action of $(\mathbb{Z}/3)^r$ with $r \geq 3$. Then $r = 3$, S is isomorphic to the Fermat cubic $X_0^3 + \dots + X_3^3 = 0$, and $(\mathbb{Z}/3)^3$ acts as the 3-torsion subgroup of the diagonal torus in $\text{PGL}_4(k)$.*

Proof: The list 3.5 gives $\ell \geq 6$. On the other hand (3.6) and Lemma 3.7 give $\ell \leq 6$. Thus S is a cubic surface in $\mathbb{P}^3$, and G is a subgroup of $\text{PGL}_4(k)$. By Lemma 3.1 we have $r = 3$, and there is a coordinate system $(X_0, \dots, X_3)$ on $\mathbb{P}^3$ such that G is the diagonal subgroup $(\mu_3)^4 / \mu_3$ of $\text{PGL}_4(k)$.

The surface S is defined by an element F of $H^0(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(3))$ which is semi-invariant with respect to the action of $(\mu_3)^4$. Under this action the space $H^0(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(3))$ is the direct sum of the invariant subspace spanned by $X_0^3, \dots, X_3^3$ and of 16 1-dimensional subspaces spanned by monomials, corresponding to 16 different characters of $(\mu_3)^4$. Since S is smooth, F must be of the form $a_0 X_0^3 + \dots + a_3 X_3^3$ with $a_i \neq 0$ for each i , hence the result. ■

Proposition 3.10.— *Let S be a del Pezzo surface and G a subgroup of $\text{Aut}(S)$ isomorphic to $(\mathbb{Z}/2)^r$ with $r \geq 4$ and $\text{rk Pic}(S)^G = 1$. Then $r = 4$, S is a quartic del Pezzo surface in $\mathbb{P}^4$ with equations $\sum_{i=0}^4 X_i^2 = \sum_{i=0}^4 \lambda_i X_i^2 = 0$ for some distinct elements $\lambda_0, \dots, \lambda_4$ of k , and G is the 2-torsion subgroup of the diagonal torus in $\text{PGL}_5(k)$.*

Proof: Once again the list 3.5 gives $\ell \geq 5$, and (3.6) and Lemma 3.7 give $\ell \leq 6$.

Suppose S is a cubic surface. Then G acts on the set of lines in S ; since 27 is odd, there must be one orbit with one element, that is, one line stable under G . This contradicts the assumption on $\text{Pic}(S)^G$.

Suppose S is an intersection of two quadrics in $\mathbb{P}^4$. Then G is a subgroup of $\text{PGL}_5(k)$; by Lemma 3.1 we have $r = 4$, and there is a coordinate system $(X_0, \dots, X_4)$ on $\mathbb{P}^4$ such that G is the diagonal subgroup $(\mu_2)^5 / \mu_2$ of $\text{PGL}_5(k)$.

The representation of $(\mu_2)^5$ on $H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(2))$ splits as

$$H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(2)) = I \oplus \sum_{i < j} k \cdot (X_i X_j),$$

where I is the invariant subspace spanned by $X_0^2, \dots, X_4^2$. The 2-dimensional subspace of quadratic forms vanishing on S must be contained in I , since otherwise it would contain some $X_i X_j$. After a change of coordinates we find the form given in the Proposition. ■

Remark 3.11.— The same method gives the subgroups of type $(\mathbb{Z}/3)^2$ of the Cremona group: besides the two conjugacy classes of subgroups of $\text{PGL}_3(k)$ described in (3.2), one gets the automorphisms groups of certain cubic surfaces and del Pezzo surfaces of degree 1. The subgroups of type $(\mathbb{Z}/2)^3$ and $(\mathbb{Z}/2)^2$ can also be described with analogous, but more tedious, methods.

4. Conjugacy classes of 2-elementary subgroups

(4.1) It follows from the theorem that the p -elementary subgroups of Cr_k of maximal order form only one conjugacy class, except in the case $p = 2$. We are going to analyze the latter case. We assume $\text{char}(k) \neq 2$ throughout this section.

For $\sigma \in \text{Cr}_k$, we denote by $\text{NF}(\sigma)$ the *normalized fixed locus* of σ , that is, the normalization of the union of the non-rational curves in $\mathbb{P}^2$ fixed by σ . The isomorphism class of $\text{NF}(\sigma)$ is an invariant of the conjugacy class of σ , and this is the basic invariant we will use to distinguish conjugacy classes.

(4.2) Let I be a finite subset of $k \setminus \{2, -2\}$; we associate to I the subgroup G_I of Cr_k generated by the involutions

$$(x, y) \mapsto (-x, y) \quad , \quad (x, y) \mapsto \left(\frac{1}{x}, y\right) \quad , \quad (x, y) \mapsto (x, -y) \quad , \quad (x, y) \mapsto \left(x, \frac{P(x)}{y}\right)$$

with $P(x) = \prod_{a \in I} (x^2 + x^{-2} - a)$. By definition this gives the conjugacy classes of subgroups of type c_1 .

Consider the Galois V_1 -covering² $\pi : \mathbb{P}_x^1 \rightarrow \mathbb{P}_u^1$ given by $u = x^2 + x^{-2}$. The normalizer $N(V_1)$ of V_1 in $\text{PGL}_2(k)$ is isomorphic to $\mathfrak{S}_4$, and the quotient

² To distinguish the two copies of $\mathbb{P}^1$ which appear we indicate the coordinate by a subscript.

$N(V_1)/V_1 \cong \mathfrak{S}_3$ acts on $\mathbb{P}_u^1$; this action preserves the branch locus $\{2, -2, \infty\}$ of π , and therefore identifies $\mathfrak{S}_3$ to the group of homographies of $\mathbb{P}_u^1$ preserving this subset. If $\gamma \in N(V_1)$, the birational transformation $(x, y) \mapsto (\gamma(x), y)$ conjugates G_I to $G_{\gamma(I)}$.

Proposition 4.3. — *The map $I \mapsto G_I$ induces a bijection between finite subsets of $k \setminus \{2, -2\}$ modulo the action of $\mathfrak{S}_3$ and conjugacy classes of subgroups of Cr_k of type c_1).*

Proof: We have already seen that the map is well-defined and surjective; it remains to prove that we can recover from G_I the finite subset $I \subset k \setminus \{2, -2\}$ modulo $\mathfrak{S}_3$.

First observe that the case $I = \emptyset$ (which gives $G_I = V_1 \times V_1 \subset \text{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$) is the only one for which $\text{NF}(\sigma) = \emptyset$ for all $\sigma \in G_I$. From now on we assume $I \neq \emptyset$.

For $\sigma \in G_I$, we have $\text{NF}(\sigma) = \emptyset$ except for $\sigma_{\pm} : (x, y) \mapsto (x, \pm \frac{P(x)}{y})$; $\text{NF}(\sigma_+)$ and $\text{NF}(\sigma_-)$ are isomorphic to the hyperelliptic curve C_I with affine equation $y^2 = P(x)$. The double covering $x : C_I \rightarrow \mathbb{P}_x^1$ is branched along $W = \pi^{-1}(I)$; the group V_1 acts freely on W .

Conversely, starting from $G_I \subset \text{Cr}_k$, we get the curve C_I , hence the branch locus $W \subset \mathbb{P}^1$ up to the action of $\text{PGL}_2(k)$. Moreover we have an action of a group $G' \cong (\mathbb{Z}/2)^2$ on $\mathbb{P}^1$ preserving W : indeed G_I acts on $C = \text{NF}(\sigma_+)$, with σ_+ acting trivially and σ_- acting as the hyperelliptic involution. We take $G' = G_I / \langle \sigma_+, \sigma_- \rangle$.

Choose a basis (e_1, e_2) of G' over $\mathbb{Z}/2$. By Lemma 2.1 there is a coordinate x on $\mathbb{P}^1$ such that e_1 acts by $x \mapsto -x$ and e_2 by $x \mapsto x^{-1}$; it is unique up to the action of V_1 . In particular the map $\pi : x \mapsto x^2 + x^{-2}$ and the subset $I = \pi(W)$ of $k \setminus \{2, -2\}$ depend only on the choice of the basis (e_1, e_2) . A change of basis is realized by an element of $N(V_1)$, so the class of $\pi(W)$ mod. $\mathfrak{S}_3$ is well-defined. Thus we recover I mod. $\mathfrak{S}_3$ from the subgroup G_I of Cr_k . ■

(4.4) We now consider subgroups of type c_2 . We first recall some very classical facts about quartic del Pezzo surfaces – one possible reference is [H], lecture 22.

A quartic del Pezzo surface $S \subset \mathbb{P}^4$ is contained in a pencil $(Q_{\lambda})_{\lambda \in \mathbb{P}^1}$ of quadrics. There are exactly 5 singular quadrics $Q_{\lambda_0}, \dots, Q_{\lambda_4}$ in this pencil; the map $S \mapsto \{\lambda_0, \dots, \lambda_4\}$ is an isomorphism from the moduli space of quartic del Pezzo surfaces onto the moduli space of 5-points subsets of $\mathbb{P}^1$ (modulo the action of PGL_2). The quadrics $Q_{\lambda_0}, \dots, Q_{\lambda_4}$ have rank 4; their singular points $p_0, \dots, p_4$ span $\mathbb{P}^4$.

The group $\text{Aut}(S)$ contains a normal, canonical subgroup G_S isomorphic to $(\mathbb{Z}/2)^4$. Indeed for $0 \leq \ell \leq 4$, there is a unique involution σ_{ℓ} of $\mathbb{P}^4$ whose fixed locus consists of p_{ℓ} and the hyperplane H_{ℓ} spanned by the points p_i for $i \neq \ell$;

these involutions span the group G_S . In more concrete terms, choose the coordinates on $\mathbb{P}^1$ so that $\lambda_0, \dots, \lambda_4 \in k$. There exists a system of coordinates on $\mathbb{P}^4$ such that the equations of Q_∞ and Q_0 are respectively

$$\sum_{i=0}^4 X_i^2 = 0 \quad \sum_{i=0}^4 \lambda_i X_i^2 = 0.$$

Then G_S is the 2-torsion subgroup of the diagonal torus in $\mathrm{PGL}_5(k)$; the involution σ_ℓ maps $(X_0, \dots, X_\ell, \dots, X_4)$ to $(X_0, \dots, -X_\ell, \dots, X_4)$. As before we view G_S as a subgroup of Cr_k , well-defined up to conjugacy.

Proposition 4.5. – a) *The subgroups of type c_1) are not conjugate to those of type c_2).*

b) *Let S and S' be two quartic del Pezzo surfaces, with S general. If $G_{S'}$ is conjugate to G_S , S' is isomorphic to S .*

In other words, the map $S \mapsto G_S$ from the moduli space of quartic del Pezzo surfaces to the set of conjugacy classes of subgroups of Cr_k is generically injective. I expect actually this map to be injective, hence a bijection onto the set of conjugacy classes of groups of type c_2); one way to prove it would be by using classical invariant theory, but the computations look rather frightening.

Proof of the Proposition : Let S be a quartic del Pezzo surface. The 5 elements σ_ℓ of G_S fix the elliptic curve $H_\ell \cap S$, while the other elements of G_S have no fixed curve. On the other hand we have seen that for a subgroup $G \subset \mathrm{Cr}_k$ of type c_1), at most 2 elements of G have a non-trivial normalized fixed locus. This is enough to prove assertion a).

(4.6) To prove b), we want to recover the surface S – or, equivalently, the 5-points subset $\{\lambda_0, \dots, \lambda_4\}$ – from the data of the elliptic curves $H_\ell \cap S$ ($0 \leq \ell \leq 4$). The curve $H_\ell \cap S$ is given in $H_\ell \cong \mathbb{P}^3$ by the equations

$$\sum_{i \neq \ell} X_i^2 = \sum_{i \neq \ell} \lambda_i X_i^2 = 0;$$

it is isomorphic to the double covering of $\mathbb{P}^1$ branched along the points λ_i for $i \neq \ell$ ([H], Prop. 22.38). The j -invariant of this curve is $j([\lambda_0, \dots, \widehat{\lambda}_\ell, \dots, \lambda_4])$, where $[\dots]$ denotes the cross-ratio and j is the rational function $j(x) = 2^8 \frac{(x^2 - x + 1)^3}{x^2(x - 1)^2}$. This function is invariant under the “cross-ratio group”, that is, the subgroup of $\mathrm{PGL}_2(k)$, isomorphic to $\mathfrak{S}_3$, generated by the involutions $z \mapsto z^{-1}$ and $z \mapsto 1 - z$. In fact the map $j : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is a Galois covering with group $\mathfrak{S}_3$.

(4.7) Let V be the complement of the diagonals in $(\mathbb{P}^1)^5$. It has two commuting actions of $\mathrm{PGL}_2(k)$ and $\mathfrak{S}_5$. We are interested in the map $J : V \rightarrow (\mathbb{P}^1)^5$

which associates to a 5-tuple $(\lambda_0, \dots, \lambda_4)$ the 5-tuple $(j_0, \dots, j_4)$ given by $j_\ell = j([\lambda_0, \dots, \widehat{\lambda_\ell}, \dots, \lambda_4])$. This map is $\mathfrak{S}_5$ -equivariant, and factors through the quotient $P := V/\mathrm{PGL}_2(k)$. Thus we have a commutative diagram

$$\begin{array}{ccc} P & \xrightarrow{J} & (\mathbb{P}^1)^5 \\ \downarrow & & \downarrow \\ P/\mathfrak{S}_5 & \xrightarrow{\bar{J}} & \mathrm{Sym}^5(\mathbb{P}^1) . \end{array}$$

The space $P/\mathfrak{S}_5$ is the moduli space of 5-points sets in $\mathbb{P}^1$, or equivalently of quartic del Pezzo surfaces; the map $\bar{J}$ associates to a quartic surface S the 5 j -invariants of the elliptic curves $H_\ell \cap S$. Our aim is to prove that $\bar{J}$ is generically injective; it suffices to prove that J is *generically injective*.

(4.8) The quotient $P = V/\mathrm{PGL}_2$ is easy to describe: any 5-tuple is equivalent to one (and only one) of the form $(\lambda, \mu, 1, 0, \infty)$. Thus we can identify P with the open subset of couples (λ, μ) in $\mathbb{P}^1 \times \mathbb{P}^1$ with $\lambda, \mu \notin \{0, 1, \infty\}$ and $\lambda \neq \mu$. Then the map J is given by

$$J(\lambda, \mu) = \left(j(\mu), j(\lambda), j\left(\frac{\lambda}{\mu}\right), j\left(\frac{\lambda-1}{\mu-1}\right), j\left(\frac{\lambda(\mu-1)}{\mu(\lambda-1)}\right) \right) .$$

Since $j^{-1}(\infty) = \{0, 1, \infty\}$ we see that J takes its value in $(\mathbb{A}^1)^5 = \mathbb{A}^5$. The strategy will be to first prove that $J : P \rightarrow \mathbb{A}^5$ is proper, then that for some $p \in P$ the tangent map $T_p(J)$ is injective and $J^{-1}(J(p)) = \{p\}$.

(4.9) Let us view J as a rational map $\mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow (\mathbb{P}^1)^5$, obtained by composition of the rational map

$$f : \mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow (\mathbb{P}^1)^5 \quad \text{given by} \quad f(\lambda, \mu) = \left(\mu, \lambda, \frac{\lambda}{\mu}, \frac{\lambda-1}{\mu-1}, \frac{\lambda(\mu-1)}{\mu(\lambda-1)} \right)$$

with the morphism $j^5 : (\mathbb{P}^1)^5 \rightarrow (\mathbb{P}^1)^5$.

The map f is well-defined except at the points $(0, 0)$, $(1, 1)$ and (∞, ∞) , and extends to a morphism $X \rightarrow (\mathbb{P}^1)^5$, where X is the surface obtained by blowing up $\mathbb{P}^1 \times \mathbb{P}^1$ along these points. Thus J extends to $\bar{J} : X \rightarrow (\mathbb{P}^1)^5$, which maps the three exceptional divisors in X into $\{\infty\} \times \{\infty\} \times (\mathbb{P}^1)^3$. Therefore $\bar{J}$ maps $X \setminus P$ to $(\mathbb{P}^1)^5 \setminus (\mathbb{A}^1)^5$, so its restriction $J : P \rightarrow \mathbb{A}^5$ is proper.

(4.10) We will now exhibit a point $p \in P$ such that $J^{-1}(J(p)) = \{p\}$. We choose an element ζ of k with $\zeta^3 = -1$; if $\mathrm{char}(k) \neq 3$ we assume moreover $\zeta \neq -1$. We

take $p = (\alpha, \zeta)$, with α general enough in k . So let $(\lambda, \mu) \neq (\alpha, \zeta)$ in P with $J(\lambda, \mu) = J(\alpha, \zeta)$. Then μ belongs to the orbit $\{\zeta, \zeta^{-1}\}$ of ζ under the cross-ratio group $\mathfrak{S}_3$.

Suppose $\mu = \zeta$, and $\text{char}(k) \neq 3$. Then we have

$$\lambda = \sigma(\alpha) \quad , \quad \frac{\lambda}{\zeta} = \tau\left(\frac{\alpha}{\zeta}\right) \quad \text{for some } \sigma, \tau \neq 1 \text{ in } \mathfrak{S}_3 .$$

This gives $\zeta^{-1}\sigma(\alpha) = \tau(\zeta^{-1}\alpha)$; if α is general enough this implies $h_\zeta^{-1}\sigma h_\zeta = \tau \in \mathfrak{S}_3$, where h_ζ is the homothety $z \mapsto \zeta z$. Since $\mathfrak{S}_3$ preserves $\{0, 1, \infty\}$ we see that σ must preserve $\{0, \infty\}$, hence fix 1; the only possibility is $\sigma(z) = z^{-1}$, which is excluded because $\zeta^2 \neq 1$.

Suppose $\mu = \zeta^{-1}$ (with no assumption on $\text{char}(k)$). We have

$$\lambda = \sigma(\alpha) \quad , \quad \zeta\lambda = \tau\left(\frac{\alpha}{\zeta}\right) \quad , \quad \zeta^{-1}(1 - \lambda) = \xi(\zeta(1 - \alpha)) \quad \text{for some } \sigma, \tau, \xi \text{ in } \mathfrak{S}_3 .$$

The first two relations give $\zeta\sigma(\alpha) = \tau(\zeta^{-1}\alpha)$, so that $h_\zeta\sigma h_\zeta = \tau \in \mathfrak{S}_3$. As before the only possibility for σ is $\sigma(z) = z^{-1}$, so that $\lambda = \alpha^{-1}$. Now the third relation becomes $\zeta^{-1}(1 - \alpha^{-1}) = \xi(\zeta(1 - \alpha))$; this cannot hold for α generic because it does not hold for $\alpha = \infty$.

(4.11) It remains to check that the tangent map to J at $p = (\alpha, \zeta)$ is injective for α general enough. The Jacobian determinant

$$\det \begin{pmatrix} \frac{\partial}{\partial \lambda} J\left(\frac{\lambda}{\mu}\right) & \frac{\partial}{\partial \mu} J\left(\frac{\lambda}{\mu}\right) \\ \frac{\partial}{\partial \lambda} J\left(\frac{\lambda-1}{\mu-1}\right) & \frac{\partial}{\partial \mu} J\left(\frac{\lambda-1}{\mu-1}\right) \end{pmatrix} = j'\left(\frac{\lambda}{\mu}\right) j'\left(\frac{\lambda-1}{\mu-1}\right) \frac{\mu - \lambda}{\mu^2(\mu - 1)^2}$$

is nonzero if $\frac{\lambda}{\mu}$ and $\frac{\lambda-1}{\mu-1}$ do not belong to $\{\zeta, \zeta^2, -1, 2, \frac{1}{2}\}$, which is certainly the case when α is general enough. This finishes the proof of the Proposition. ■

REFERENCES

- [B-B] L. BAYLE, A. BEAUVILLE: *Birational involutions of $\mathbb{P}^2$* . Kodaira's issue, Asian J. Math. **4** (2000) 11–17.
- [B-Bl] A. BEAUVILLE, J. BLANC: *On Cremona transformations of prime order*. C. R. Math. Acad. Sci. Paris **339** (2004), no. 4, 257–259.
- [B] N. BOURBAKI: *Groupes et algèbres de Lie*, Chap. VI. Hermann, Paris (1968).
- [Bo] A. BOREL: *Sous-groupes commutatifs et torsion des groupes de Lie compacts connexes*. Tôhoku Math. J. (2) **13** (1961), 216–240.

- [D1] M. DEMAZURE: *Sous-groupes algébriques de rang maximum du groupe de Cremona*. Ann. Sci. École Norm. Sup. (4) **3** (1970), 507–588.
- [D2] M. DEMAZURE: *Surfaces de del Pezzo*. Séminaire sur les Singularités des Surfaces. LNM **777**, 21–69. Springer, Berlin (1980).
- [Do] I. DOLGACHEV: *Weyl groups and Cremona transformations*. Singularities I, 283–294, Proc. Sympos. Pure Math. **40**, AMS, Providence (1983).
- [dF] T. DE FERNEX: *On planar Cremona maps of prime order*. Nagoya Math. J. **174** (2004), 1–28.
- [dF-E] T. DE FERNEX, L. EIN: *Resolution of indeterminacy of pairs*. Algebraic geometry, 165–177, de Gruyter, Berlin (2002).
- [H] J. HARRIS: *Algebraic geometry, a first course*. Graduate Texts in Mathematics, **133**. Springer-Verlag, New York (1995).
- [K] S. KANTOR: *Theorie der endlichen Gruppen von eindeutigen Transformationen in der Ebene*. Mayer & Müller, Berlin (1895).
- [M] Y. MANIN: *Rational surfaces over perfect fields*, II. Math. USSR – Sbornik **1** (1967), 141–168.
- [W] A. WIMAN: *Zur Theorie der endlichen Gruppen von birationalen Transformationen in der Ebene*. Math. Ann. **48** (1896), 195–240.
- [Z] D.-Q. ZHANG: *Automorphisms of finite order on rational surfaces*. J. Algebra **238** (2001), no. 2, 560–589.

Arnaud BEAUVILLE
 Institut Universitaire de France
 &
 Laboratoire J.-A. Dieudonné
 UMR 6621 du CNRS
 UNIVERSITÉ DE NICE
 Parc Valrose
 F-06108 NICE Cedex 02